



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention  
of the grant of the patent:  
**26.06.2002 Bulletin 2002/26**

(51) Int Cl.7: **A61K 7/06, C12N 5/06,**  
**C12N 5/00**

(21) Application number: **98908329.0**

(86) International application number:  
**PCT/NL98/00129**

(22) Date of filing: **05.03.1998**

(87) International publication number:  
**WO 98/47471 (29.10.1998 Gazette 1998/43)**

(54) **METHOD FOR THE PROPAGATION OF HAIR**  
**VERFAHREN ZUR VERMEHRUNG VON HAAREN**  
**PROCEDE DE REPRODUCTION DES CHEVEUX**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC**  
**NL PT SE**

(30) Priority: **05.03.1997 NL 1005445**

(43) Date of publication of application:  
**19.01.2000 Bulletin 2000/03**

(73) Proprietor: **Gho Holding B.V.**  
**6216 BX Maastricht (NL)**

(72) Inventor: **GHO, Conradus, Ghosal**  
**NL-6241 JL Bunde (NL)**

(74) Representative: **de Bruijn, Leendert C. et al**  
**Nederlandsch Octrooibureau**  
**P.O. Box 29720**  
**2502 LS Den Haag (NL)**

(56) References cited:  
**EP-A- 0 236 014** **EP-A- 0 682 107**  
**WO-A-92/07877** **WO-A-95/01423**  
**WO-A-96/32961**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

## Description

**[0001]** The present invention firstly relates to a method for the reproduction of hair. The invention also relates to a method for culturing hair follicle cells from anagenetic hairs in the anagen phase, to a medium that is highly suitable for culturing hair follicle cells, to a method for the preparation of such a medium and to a precursor medium.

**[0002]** A method for the reproduction of hair is disclosed in European Patent Application 0 236 014, in which epidermal follicle cells of the desired hair type are removed from the scalp skin of a patient. The epidermal follicle cells are then cultured in a culture medium which preferably contains growth factors. In a subsequent step, an opening is made in the epidermis of the patient's scalp and, via said opening, the cultured epidermal follicle cells are introduced into the dermis next to the epidermis. The disadvantage of this method is that it comprises an invasive method and that the cells are not placed directionally, as a result of which many cells are necessary and the probability of regeneration of hair is much lower. In this method, use is also not made of autologous (cultured) CD34<sup>+</sup> cells.

**[0003]** The essential growth structures of hair are the so-called hair follicles, which are present in the skin. The hair follicle cells or keratinocytes reproduce from these hair follicles and during their path to the skin surface the cytoplasm of said cells is converted by a large number of complex processes into the tough and resilient material which is known as hair. The growth cycle of hair can be subdivided into three phases: the anagen phase or growth phase, the catagen phase or transitory phase and the telogen phase or dying phase. The hair follicle is unique because of the cyclic nature of hair formation and hair growth. Specifically, the hair follicle is the only part of the body to have a growth nucleus, from which new hairs can be produced after removal of the old hair.

**[0004]** It is known that hair follicle cells from plucked human hair can be cultured. It is also known that it is possible using such cultured cells to form a differentiated epidermis or a fully developed epidermis, both in vitro and in vivo. Cultured hair follicle cells from mice can stimulate hair growth when said cells are implanted into test animals. However, to date it has not proved possible to achieve new hair growth in humans in those locations where there is undesirably no (longer any) hair with the aid of cultured autologous hair follicle cells.

**[0005]** People usually find baldness to be undesirable from the cosmetic and aesthetic standpoint. However, baldness frequently occurs and it is a known phenomenon that in particular men also become balder as they get older. This form of baldness is known as alopecia androgenetica. To date it is not precisely known why certain parts of the scalp are susceptible to this alopecia androgenetica and other parts are not. However, with women as well it regularly occurs that the hair becomes thinner and even threatens to largely disappear. For

women in particular this is highly undesirable from the cosmetic and aesthetic standpoint.

**[0006]** A known technique for combatting baldness is to transplant hair. With this procedure hair from a donor region covered in hair, which frequently is located on the back of the head, is removed, including the skin, and cut into small pieces, which usually then have only one to three hairs. These pieces are then implanted in the bald region (receptor region). A major disadvantage of this method is that it is at the cost of the donor region. Specifically, hair is removed from this region and this hair does not return again. The transplantation technique therefore offers limited possibilities.

**[0007]** The aim of the present invention is to provide a technique with the aid of which bald patches can be provided with hair again, but which does not have the disadvantages of hair transplantation which have been outlined above. A further aim or the invention is to provide a method with which new hair growth on bald patches can be achieved in humans with the aid of cultured hair follicle cells. A further aim of the invention is to provide a method which is relatively simple to carry out.

**[0008]** Said aims are achieved with the method according to the invention, wherein hair is, as it were, reproduced. Specifically, according to the invention the hair is removed from a donor region in such a way that new hairs come back in its place, whilst new hair follicle cells are cultured from the hairs removed, from which cells, in turn, new hair can form.

**[0009]** The invention therefore relates to a cosmetic method for the reproduction of hair, which method comprises the following steps:

- (1) removal of hair in the anagen phase from one or more donor regions in such a way that the bulb characteristic of hair in the anagen phase is still attached to the hair removed.
- (2) culture of hair follicle cells from the hair removed, under circumstances such that the hair follicle cells are able to multiply, and
- (3) implantation of the cultured hair follicle cells in the receptor regions.

wherein:

- the hair in the anagen phase is removed by plucking hair out of the donor region or the donor regions, followed by selection of suitable hairs in the anagen phase,
- the hair follicle cells are cultured in a culture medium which is optionally supplemented with (a) at least one human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells, or (b) one or more extracts of the human mast cell line(s) and/or of the CD34<sup>+</sup> cells and/or (c) growth-stimulating agents, and
- the cultured follicular hair cells and/or the autologous (cultured) CD34<sup>+</sup> cells are implanted in the pores of the receptor region.

[0010] According to the invention, only one hair follicle cell or one autologous (cultured) CD34<sup>+</sup> cell or a combination of these two cells can be introduced although a plurality of cells is usually mentioned in the description.

[0011] The method according to the invention has the major advantage that hair growth can be achieved again on bald patches without this being at the expense of the donor region.

[0012] Step (1) of the method according to the invention comprises the removal of hair in the anagen phase from a donor region where such hairs are located. As explained above, hair growth comprises three phases: an anagenous, a catagenous and a telogenous phase. Only hairs which are in the anagenous phase are suitable for the method according to the invention. Compared with hairs in the catagen and telogen phases, such hairs in the anagen phase are characterized in that they have a frequently pigmented bulb of a shape characteristic for hairs in the anagen phase at the bottom of the hair. This is generally known and to an experienced eye a hair in the anagen phase is therefore also immediately recognizable from the shape of the bulb. In the case of doubt the use of a microscope can provide a definite answer. The removal of hairs in the anagen phase can therefore be effected in various ways, as long as the bulb characteristic of hairs in the anagen phase is still attached to the hair removed.

[0013] The number of hairs in the anagen phase needed to culture a suitable quantity of hair follicle cells is hardly subject to specific limits since in principle an unlimited number of hair follicle cells can be cultured from a single hair. Specifically, subcultures can, in turn, be cultured from the first culture and, in turn, several cultures can be cultured from every subculture. In practice, 3 to 10 hairs in the anagen phase will be ample. The hairs which ultimately are formed in the receptor region from the cultured hair follicle cells assume the characteristics of the hairs from the donor region from which the hair follicle cells were cultured. Therefore, a region which is not susceptible to alopecia androgenetica is taken as a suitable donor region. However, this is not essential.

[0014] Since the difference between hairs in the various growth stages can be seen most clearly once the hairs have been removed, the hairs from the donor region are removed by plucking hair out of the donor region and selecting the suitable hairs in the anagen phase. Tweezers are, for example, very suitable for plucking hair from the donor region.

[0015] In step (2) of the method according to the invention the hairs are cultured from the hair removed, under conditions such that the hair follicle cells are able to multiply. In principle, the hair follicle cells can be cultured from the hairs by the known method. Culture media for this purpose are available commercially and any associated growth supplements are freely available. Such culture media are also termed serum-free keratinocyte culture media and usually contain essential and non-es-

sential amino acids, vitamins, trace elements, organic constituents and inorganic salts. Such culture media can also be combined with growth supplements, which contain growth factors, hormones, antibiotics and tissue extracts, such as, for example, BPE (Bovine Pituitary Extract), insulin, hydrocortisone, HEGF (Human Epidermal Growth Factor), TGF (Transformal Growth Factor), L-cysteine, L-leucine and gentamicin. According to the invention, particularly good results are obtained if, in step (2) of the method according to the invention, use is made of a culture medium which, in addition to the known culture media described above, is optionally supplemented with one or more growth supplements, and/or is supplemented with one human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells (CD34 positive cells) or one or more extracts of the human mast cell line(s) and/or of the CD34<sup>+</sup> cells and/or growth-stimulating agents. Preferably, only cultured CD34<sup>+</sup> cells are used because these cells are native to the body. Human mast cells are known per se. They contain various growth factors and are produced and consumed in various different sites in the body. The extracts of the human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth-stimulating agents contain growth factors. However, to date such extracts of the human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth-stimulating agents have not been used in media for culturing hair follicle cells. A human mast cell line from which the extracts can be used, is HMC-1 (Human Mast Cell line 1) or a cell line derived therefrom, such as the subclones 5C6 and KUB12. Mast cell lines such as HMC-1 and subclones thereof are commercially available. According to the invention, instead of mast cells, autologous (cultured) CD34<sup>+</sup> cells or a combination of two or more of these cell types can be used.

[0016] The extracts of the human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth-stimulating agents can very suitably be introduced into the medium by adding at least one degranulating agent to the serum-free keratinocyte culture medium containing the human mast cell line, a subclone thereof and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth agents. An inorganic salt can also be added to optimise the action of the degranulating agent. A suitable salt in this connection is CaCl<sub>2</sub>. The degranulating agent then, as it were, cuts the mast cells and/or the autologous (cultured) CD34<sup>+</sup> cells into smaller pieces (degranulation), as a result of which the growth factors are released into the medium. After degranulation, the residues of the mast cells and/or the autologous (cultured) CD34<sup>+</sup> cells can be removed, for example by means of centrifuging. The resultant, conditioned medium can then be used to culture the hair follicle cells.

[0017] Degranulating agents are known per se and can roughly be divided into three groups:

- hormones, such as oestrogens and bradykinin,
- neurotransmitters and antigens, such as apitoxin.

- Substance P and Ig E, and synthetic agents, such as Compound 48/80 and ionophore A23187. Those degranulating agents which are capable of degranulating the mast cells optionally used are suitable for use in the method according to the invention.

**[0018]** An alternative is first separately to prepare the extracts of the human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth agents and to add these extracts as such to the serum-free keratinocyte culture medium.

**[0019]** The conditions under which the hair follicle cells are cultured in step (2) can vary substantially and are dependent on, inter alia, the medium used. However, it has been found that the culturing of the hair follicle cells at a temperature of 30-40°C, preferably in an incubator so that the temperature can be kept constant, in an atmosphere which is saturated with water and which contains 3-10% by volume CO<sub>2</sub> in addition to the other customary constituents of air in the customary quantities gives good results. However, these conditions are in no way restrictive and other conditions can therefore also be used. The culture period varies per culture, usually from 1 hour to 40 days. In order to obtain a good result, it is preferable to replace the medium by fresh medium every 2 to 5 days.

**[0020]** Step (2) of the method according to the invention can be carried out in a single culture step, but can also be carried out in several substeps by culturing subcultures from the initial culture or cultures. This can be carried out by taking a sufficiently large number of hair follicle cells from the main culture when the latter contains sufficient hair follicle cells. The hair follicle cells from the main culture are cultured in turn, preferably - but not necessarily - in the same culture medium as that in which the original main culture was cultured. In the course of time it is possible, if desired, to culture yet further cultures from said subcultures, etc. etc.

**[0021]** After the hair follicle cells have been cultured they must be detached from the substrate on which they have been grown. This can be carried out by the known methods, for example by making use of a trypsin solution (0.1-0.5% by weight in water), optionally in combination with an EDTA solution (0.01-0.05% by weight) in PBS (Phosphate Buffered Saline). Such methods are known per se. Before the cultured hair follicle cells are used in step (3), the mast cells and/or autologous (cultured) CD34<sup>+</sup> cells can, if desired, be removed.

**[0022]** In step (3) of the method according to the invention, the cultured hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells are implanted in the skin in the receptor region. It has been found that very good results are obtained when the cultured hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells are implanted directly into the pores of the receptor region. If the pores are difficult to see, which can with a pale skin colour in particular, the pores can be made more readily visible by

applying a (usually yellow-coloured) liquid to the skin in the receptor region, which becomes darker in colour in the positions where the pores are located as a result of the secretion of the perspiration from the sweat glands present in the pores. An example of such a liquid is a 1-10% solution of o-phthalaldehyde in, for example xylene or ethyl ether, as described by L. Juhlin and W. B. Suelley in Nature, 28 January 1967, page 408.

**[0023]** Preferably a quantity of a suspension of cultured hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells is introduced into each pore such that said quantity contains sufficient hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells to allow a hair follicle to develop from which a hair is able to grow. The number of cultured hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells needed to cause a hair follicle to form is highly dependent on the degree of differentiation or the growth potential of the cultured cells. Naturally, fewer cultured cells will be needed when the cultured hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells have a relatively high degree of differentiation than when the cultured cells have a relatively low degree of differentiation. There are a large number of factors which influence the degree of differentiation of the cultured hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells. For example, the medium used and the culture time are factors which play a role. Preferably, a suspension of the cultured hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells in the medium in which said cells were cultured is used. The suspension preferably has a hair follicle cell concentration which is such that 0.01-5 µl, preferably 0.1 to 0.5 µl, of suspension per pore can suffice. A small quantity such as this is preferably injected into the pore with the aid of a metering device which is provided with a hollow needle, preferably an 18 to 40 Gauge (G) needle. The metering device used can be, for example, an electronic metering system for a pipette, a 18-40 G needle being fitted on the end of the metering hose or of the pipette. In the light of the average size of the pores in, in particular, the scalp, it is preferable to use a hollow 34 G needle. However, smaller or larger needles can also be used on other parts of the body. It is also possible and preferable for the suspension to be injected with an (electronic) repeating injection-metering apparatus such as an insulin pen. In that case, a needle specially developed for the purpose, that is to say a "follicle-conducting needle", is also used to introduce the cells at the desired position and to damage the skin as little as possible. A follicle-conducting needle is a needle which is flexible and minuscule and also which the hair follicle can follow. During the injection of the suspension, substances can also be introduced at the same time which keep the cells alive for longer. Examples of such substances are Minoxidil and other growth-stimulating agents.

**[0024]** The invention also relates to a method for culturing hair follicle cells from hairs in the anagen phase, one or more hairs in the anagen phase being placed in

a medium which is supplemented with one suitable serum-free keratinocyte culture medium and, optionally, supplemented with one human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells or one or more extracts of the human mast cell line(s) and/or of the CD34<sup>+</sup> cells and/or growth-stimulating agents. The way in which such a medium can be obtained has already been described above.

**[0025]** As far as the preferences with regard to the various constituents of the medium to be used and to the culturing conditions are concerned, these are the same as those for the medium and the conditions employed in step (2) of the method already described above. Therefore, it is preferable to use one mast cell line, a subclone thereof and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth-stimulating agents. Suitable conditions comprise a culture temperature of 30-40°C in an atmosphere which is saturated with water and contains 3-10% by volume of CO<sub>2</sub>.

**[0026]** The invention also relates to a medium for culturing hair follicle cells from hairs in the anagen phase and to a method for the preparation of said medium. The medium according to the invention comprises at least one suitable serum-free keratinocyte culture medium, optionally supplemented with one human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells or extracts of the human mast cell line(s) and/or of the CD34<sup>+</sup> cells and/or growth-stimulating agents. The special feature of such a medium is, in particular, that it contains growth factors originating from one or more human mast cell lines and/or autologous (cultured) CD34<sup>+</sup> cells.

**[0027]** A preferred medium comprises a serum-free keratinocyte culture medium and extracts of the human mast cell line HMC-1 or a cell line derived therefrom, such as 5C6 or KU812 or autologous (cultured) CD34<sup>+</sup> cells. Further details on the constituents of the medium according to the invention have already been given above.

**[0028]** The method for the preparation of said medium comprises the following steps:

- (a) addition of at least one mast cell line, subclone thereof and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth-stimulating agents and optionally one degranulating agent, optionally in combination with an organic salt, such as, for example, CaCl<sub>2</sub>, to a serum-free keratinocyte culture medium;
- (b) degranulation of the mast cell line(s), subclones thereof and/or autologous (cultured) CD34<sup>+</sup> cells, so that the growth factors are released into the medium; and
- (c) optionally removal of the mast cells and/or autologous (cultured) CD34<sup>+</sup> cells from the medium, for example by centrifuging, after which the (conditioned) medium is obtained.

**[0029]** According to the invention, the mast cells and/or the CD34<sup>+</sup> cells and/or the residues thereof are re-

moved from the medium.

**[0030]** The conditions under which this method can be carried out are not particularly critical and will be obvious to any person skilled in the art. The degranulating agent used in step (a) is preferably Compound 48/80 in combination with CaCl<sub>2</sub>.

**[0031]** Finally, the invention relates to a precursor medium for culturing hair follicle cells from hairs in the anagen phase which medium comprises at least one suitable serum-free keratinocyte culture medium and which is optionally supplemented with one human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells or one or more extracts of the human mast cell line(s) and/or of the CD34<sup>+</sup> cells and/or growth-stimulating agents and/or at least one degranulating agent. Said precursor medium can be used as such for preparing therefrom a suitably conditioned medium for culturing hair follicle cells from hairs in the anagen phase. This can be performed simply by carrying out the degranulation if both a human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells is/are present as well as degranulating agent. If one of the two is absent, that is to say the mast cells and/or the CD34<sup>+</sup> cells and the degranulating agent, respectively, the missing component must be first added before degranulation of the mast cells can be carried out.

#### Claims

1. Cosmetic method for the reproduction of hair, which method comprises the following steps:

- (1) removal of hair in the anagen phase from one or more donor regions in such a way that the bulb characteristic of hair in the anagen phase is still attached to the hair removed,
- (2) culture of hair follicle cells from the hair removed, under circumstances such that the hair follicle cells are able to multiply, and
- (3) implantation of the cultured hair follicle cells in the receptor regions,

wherein:

- the hair in the anagen phase is removed by plucking hair out of the donor region or the donor regions, followed by selection of suitable hairs in the anagen phase,
- the hair follicle cells are cultured in a culture medium which is optionally supplemented with (a) at least one human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells, or (b) one or more extracts of the human mast cell line(s) and/or of the CD34<sup>+</sup> cells and/or (c) growth-stimulating agents, and
- the cultured follicular hair cells and/or the autologous (cultured) CD34<sup>+</sup> cells are implanted in the pores of the receptor region.

2. Method according to Claim 1, wherein the human mast cell line is one mast cell line, a subclone thereof and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth-stimulating agents.
3. Method according to Claim 2, wherein the human mast cell line is one human mast cell line and/or an autologous (cultured) CD34<sup>+</sup> cell line.
4. Method according to one of the preceding claims, wherein the hair follicle cells are cultured at a temperature of 30-40°C in an atmosphere which is saturated with water and contains 3-10% by volume of CO<sub>2</sub>.
5. Method according to Claim 1, wherein a quantity of a suspension of cultured hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells is introduced into each pore such that said quantity contains sufficient hair follicle cells and/or autologous (cultured) CD34<sup>+</sup> cells to allow a hair follicle to develop.
6. Method according to Claim 5, wherein the quantity of suspension introduced per pore is 0.01-5 µl.
7. Method according to Claim 6, wherein the quantity is 0.1-0.5 µl.
8. Method according to one of Claims 5-7, wherein the suspension is introduced into each pore individually via a metering pipette which is provided with a hollow needle, preferably an 18 to 40 Gauge needle.
9. Method according to one of Claims 5-7, wherein the suspension is introduced via a repeating injection-metering apparatus.
10. Method according to Claim 9, wherein a follicle-conducting needle is used.
11. Use of hair follicle cells cultured from hair in the anagen phase as active component in the preparation of an injection suspension for implantation in the pores of the receptor region for the reproduction of hair, said hair being removed from one of more donor regions in such a way that the bulb characteristic of hair in the anagen phase is still attached to said hair and said hair being removed by plucking hair out of said donor region or donor regions followed by selecting suitable hairs in the anagen phase, said hair follicle cells being cultured from said suitable hairs in the anagen phase under circumstances such that the hair follicle cells are able to multiply and said hair follicle cells being cultured in a culture medium, said culture medium being optionally supplemented with (a) at least one human mast cell line and/or autologous (cultured) CD34<sup>+</sup> cells, or (b) one or more extracts of the human mast

cell line(s) and/or of the CD34<sup>+</sup> cells and/or (c) growth-stimulating agents.

12. Use according to claim 11, wherein the human mast cell line is one mast cell line, a subclone thereof and/or autologous (cultured) CD34<sup>+</sup> cells and/or growth-stimulating agents.
13. Use according to claim 12, wherein the human mast cell line is one human mast cell line and/or an autologous (cultured) CD34<sup>+</sup> cell line.
14. Use according to anyone of claims 11-13, wherein the hair follicle cells are cultured at a temperature of 30-40° in an atmosphere which is saturated with water and contains 3-10% by volume of CO<sub>2</sub>.

#### Patentansprüche

1. Kosmetisches Verfahren zur Reproduktion von Haar, wobei das Verfahren die folgenden Schritte umfaßt:
  - (1) Entfernen von Haar in der anagenen Phase aus einem oder aus mehreren Spenderbereichen, so daß die Bulbus-Charakteristik von Haar in der anagenen Phase noch mit dem entfernten Haar verbunden ist,
  - (2) Züchten von Haarfollikelzellen vom entfernten Haar unter Bedingungen, unter denen die Haarfollikelzellen dazu in der Lage sind, sich zu vermehren, und
  - (3) Einpflanzen der gezüchteten Haarfollikelzellen in die Empfängerbereiche, worin
    - das Haar in der anagenen Phase durch Herausreißen aus dem Spenderbereich oder den Spenderbereichen entfernt wird, gefolgt von der Auswahl von geeigneten Haaren in der anagenen Phase,
    - die Haarfollikelzellen in einer Nährlösung gezüchtet werden, die wahlweise mit (a) mindestens einer menschlichen Mastzelllinie und/oder autologen (gezüchteten) CD34<sup>+</sup>-Zellen oder (b) einem oder mehreren Extrakten der menschlichen Mastzelllinie(n) und/oder der CD34<sup>+</sup>-Zellen und/oder (c) wachstumsstimulierenden Mitteln ergänzt ist, und
    - die gezüchteten Follikularhaarzellen und/oder die autologen (gezüchteten) CD34<sup>+</sup>-Zellen in die Poren des Empfängerbereichs eingepflanzt werden.

2. Verfahren nach Anspruch 1, worin die menschliche Mastzelllinie eine Mastzelllinie, ein Subklon davon und/oder autologe (gezüchtete) CD34<sup>+</sup>-Zellen und/oder wachstumsstimulierende Mittel darstellt. 5
3. Verfahren nach Anspruch 2, worin die menschliche Mastzelllinie eine Mastzelllinie und/oder eine autologe (gezüchtete) CD34<sup>+</sup>-Zelllinie ist.
4. Verfahren nach einem der vorstehenden Ansprüche, worin die Haarfollikelzellen bei einer Temperatur von 30 - 40 °C in einer Atmosphäre gezüchtet werden, die mit Wasser gesättigt ist und 3 - 10 Volumenprozent CO<sub>2</sub> enthält. 10
5. Verfahren nach Anspruch 1, worin eine Menge einer Suspension von gezüchteten Haarfollikelzellen und/oder autologen (gezüchteten) CD34<sup>+</sup>-Zellen in jede Pore eingeführt wird, so daß die Menge ausreichend Haarfollikelzellen und/oder autologe (gezüchtete) CD34<sup>+</sup>-Zellen enthält, um die Entwicklung eines Haarfollikels zu ermöglichen. 15
6. Verfahren nach Anspruch 5, worin die Menge der Suspension, die pro Pore eingeführt wird, 0,01 - 5 µl beträgt. 20
7. Verfahren nach Anspruch 6, worin die Menge 0,1 - 0,5 µl beträgt. 25
8. Verfahren nach einem der Ansprüche 5 - 7, worin die Suspension in jede Pore einzeln über eine Meßpipette eingeführt wird, die mit einer hohlen Nadel, vorzugsweise einer 18 bis 40 Gauge-Nadel, ausgestattet ist. 30
9. Verfahren nach einem der Ansprüche 5 - 7, worin die Suspension mit Hilfe einer Vorrichtung zur wiederholten Einspritz-Messung eingeführt wird. 35
10. Verfahren nach Anspruch 9, worin eine follikelführende Nadel verwendet wird. 40
11. Verwendung von Haarfollikelzellen, die aus Haar in der anagenen Phase als aktiver Bestandteil bei der Zubereitung einer Einspritzsuspension zur Einpflanzung in die Poren des Empfängerbereichs für die Reproduktion von Haar gezüchtet werden, wobei das Haar aus einem oder mehreren Spenderbereichen so entfernt wird, daß die Bulbus-Charakteristik von Haar in der anagenen Phase noch mit dem Haar verbunden ist, und wobei das Haar durch Herausreißen aus dem Spenderbereich oder den Spenderbereichen entfernt wird, gefolgt von der Auswahl von geeigneten Haaren in der anagenen Phase, wobei die Haarfollikelzellen von den geeigneten Haaren in der anagenen Phase unter Bedingungen gezüchtet werden, unter denen die Haar- 45

follikelzellen dazu in der Lage sind, sich zu vermehren, und wobei die Haarfollikelzellen in einer Nährlösung gezüchtet werden, wobei die Nährlösung wahlweise mit (a) mindestens einer menschlichen Mastzelllinie und/oder autologen (gezüchteten) CD34<sup>+</sup>-Zellen, oder (b) einem oder mehreren Extrakten der menschlichen Mastzelllinie(n) und/oder der CD34<sup>+</sup>-Zellen und/oder (c) wachstumsstimulierenden Mitteln ergänzt ist.

12. Verwendung nach Anspruch 11, worin die menschliche Mastzelllinie eine Mastzelllinie ist, ein Subklon davon und/oder autologe (gezüchtete) CD34<sup>+</sup>-Zellen und/oder wachstumsstimulierende Mittel. 15

13. Verwendung nach Anspruch 12, worin die menschliche Mastzellenlinie eine menschliche Mastzellenlinie und/oder eine autologe (gezüchtete) CD34<sup>+</sup>-Zelllinie ist. 20

14. Verwendung nach irgendeinem der Ansprüche 11 - 13, worin die Haarfollikelzellen bei einer Temperatur von 30 - 40° in einer Atmosphäre gezüchtet werden, die mit Wasser gesättigt ist und 3 - 10 Volumenprozent CO<sub>2</sub> enthält. 25

#### Revendications

1. Méthode cosmétique pour la reproduction de cheveux, ladite méthode comportant les étapes suivantes : 30

(1) prélèvement de cheveux en phase anagène dans une ou plusieurs régions donneuses de manière à ce que le bulbe de cheveu 35

caractéristique de cheveux en phase anagène est encore attaché au cheveu enlevé.

(2) culture de cellules de follicules de cheveux à partir des cheveux prélevés dans des circonstances propres à permettre aux cellules de follicules de se multiplier, et 40

(3) implantation des cellules cultivées de follicules de cheveux dans les régions réceptrices, caractérisée en ce que 45

- le cheveu en phase anagène est prélevé en arrachant les cheveux des régions donneuses, des cheveux appropriés étant ensuite sélectionnés en phase anagène, 50

- les cellules de follicules de cheveux sont cultivées dans un milieu de culture complété en option par (a) au moins une lignée mastocyte humaine et/ou des cellules autologues (cultivées) CD34<sup>+</sup> ou (b) un ou 55

- plusieurs extraits de la (des) lignée(s) mastocyte(s) humaine(s) et/ou des cellules CD34<sup>+</sup> et/ou (c) des agents stimulateurs de croissance, et
- les cellules de culture de follicules de cheveux et/ou les cellules autologues (cultivées) CD34<sup>+</sup> sont implantées dans les pores de la région réceptrice.
2. Méthode selon la revendication 1, **caractérisée en ce que** la lignée mastocyte humaine est une lignée mastocyte, un sous-clone de celle-ci et/ou des cellules autologues (cultivées) CD34<sup>+</sup> et/ou des agents stimulateurs de croissance.
  3. Méthode selon la revendication 2, **caractérisée en ce que** la lignée mastocyte humaine est une lignée mastocyte humaine et/ou une lignée cellulaire autologue (cultivée) CD34<sup>+</sup>.
  4. Méthode selon une des revendications précédentes, **caractérisée en ce que** les cellules de follicules de cheveux sont cultivées à une température de 30 à 40 °C dans une atmosphère saturée en eau et contenant 3 à 10 % en volume de CO<sub>2</sub>.
  5. Méthode selon la revendication 1, **caractérisée en ce qu'**une quantité de suspension de cellules cultivées de follicules de cheveux et/ou de cellules autologues (cultivées) CD34<sup>+</sup> est introduite dans chaque pore de manière à ce que ladite quantité contienne suffisamment de cellules de follicules de cheveux et/ou de cellules autologues (cultivées) CD34<sup>+</sup> pour permettre à un follicule de cheveux de se développer.
  6. Méthode selon la revendication 5, **caractérisée en ce que** la quantité de suspension introduite par pore est de 0,01 à 5 µl.
  7. Méthode selon la revendication 6, **caractérisée en ce que** la quantité est de 0,1 à 0,5 µl.
  8. Méthode selon une des revendications 5 à 7, **caractérisée en ce que** la suspension est introduite dans chaque pore individuellement par une pipette doseuse munie d'une aiguille creuse, de préférence une aiguille de 18 à 40 Gauge.
  9. Méthode selon une des revendications 5 à 7, **caractérisée en ce que** la suspension est introduite par l'intermédiaire d'un appareil d'injection dosée à répétition.
  10. Méthode selon la revendication 9, **caractérisée en ce qu'**une aiguille conductrice de follicules est utilisée.
  11. Utilisation de cellules de follicules de cheveux cultivées à partir de cheveux en phase anagène en tant que composant actif dans la préparation d'une suspension à injecter pour implantation dans les pores de la région réceptrice pour la reproduction de cheveux, lesdits cheveux étant prélevés dans une ou plusieurs régions donneuses de manière à ce que les bulbes desdits cheveux caractéristiques en phase anagène sont encore attachés au cheveux et lesdits cheveux étant prélevés en arrachant des cheveux de ladite ou desdites région(s) donneuse(s), des cheveux appropriés étant ensuite sélectionnés en phase anagène, les cellules de follicules desdits cheveux étant cultivées à partir desdits cheveux appropriés en phase anagène dans des circonstances telles que les cellules de follicules de cheveux puissent se multiplier et les cellules de follicules desdits cheveux étant cultivées dans un milieu de culture, ledit milieu de culture étant complété en option par (a) au moins une lignée mastocyte humaine et/ou des cellules autologues (cultivées) CD34<sup>+</sup>, ou (b) un ou plusieurs extraits de la(des) lignée(s) mastocyte(s) humaine(s) et/ou des cellules CD34<sup>+</sup> et/ou (c) des agents stimulateurs de croissance.
  12. Utilisation selon la revendication 11, **caractérisée en ce que** la lignée mastocyte humaine est une lignée mastocyte, un sous-clone de celle-ci et/ou des cellules autologues (cultivées) CD34<sup>+</sup> et/ou des agents stimulateurs de croissance.
  13. Utilisation selon la revendication 12, **caractérisée en ce que** la lignée mastocyte humaine est une lignée mastocyte et/ou une lignée cellulaire autologue (cultivée) CD34<sup>+</sup>.
  14. Utilisation selon n'importe laquelle des revendications 11 à 13, **caractérisée en ce que** les cellules de follicules de cheveux sont cultivées à une température de 30 à 40° dans une atmosphère saturée en eau et contenant 3 à 10 % en volume de CO<sub>2</sub>.